

A large crayfish is shown from a top-down perspective against a solid blue background. The crayfish is oriented horizontally, with its head to the left and tail to the right. Its two large claws are prominent, and its long antennae extend forward. The body is segmented and has a dark, mottled appearance.

“THE COMMERCIAL YABBY FARMER”

A 250 PAGE BOOK ON SEMI INTENSIVE  
YABBY FARMING

by **ROBERT B McCORMACK**

# POND SIZE

- **Pond Size is important, small ponds under 1000mtrs square surface area produce small yabbies, if you want big yabbies you need big ponds. Ponds over 5000m<sup>2</sup> are too big and unmanageable for semi intensive farming.**
- **You need ponds that are gravity drainable and large ponds just take too long to drain and effectively harvest any yabbies left in the pond. Drainable ponds are a must to effectively control yabby populations and predators.**

# SHALLOW

- Yabbies only use floor space, not a water column so shallow is good.
- Yabbies are cold blooded animals, the warmer the water the faster their metabolism goes and the faster they grow. Shallow water heats rapidly in the sun.
- Shallow water does not stratify (reducing oxygen at the bottom).
- 600mm is the minimum depth or predation from birds is a serious concern.
- 900mm the preferred depth

# TURBID

- Essential to have silts/clays in suspension to give you turbid water.
- Turbid water blocks sunlight penetration into the water column reducing algal populations and water quality fluctuations.
- Sunlight impacts on clay particles in suspension which heats your water.
- Yabbies feed during the day in turbid water so grow fast. In clear water hide during day and feed only at night.
- Harvesting can be done during the day in turbid water. In clear water only at night as yabbies will hide during the day.

# TURBID (continued)

- **Burrowing is essential in clear water for yabbies to hide, in turbid water not required.**
- **Harder for predators to find yabbies in turbid water.**
- **Only high densities of yabbies will be achieved in turbid water. In clear water yabbies can see each other and one raises a claw then the other raises a claw and a fight will break out. This aggression is contagious and the whole population becomes aggressive and cannibalistic. In turbid water the density of yabbies can increase 10 fold if food is available as the yabbies can not see each other and must feel each other up with their antenna. Once they have felt each other up they are good mates and get along just fine.**

# CONSTANT WATER LEVEL

- **It is essential that little variation in water levels occur in yabby ponds.**
- **Yabbies are creatures that have evolved in Australia which has extremes in climate. In the wild if water levels drop then yabbies start burrowing. In your ponds if water levels start to drop the yabbies will burrow. You do not want them wasting energy digging holes you want them sitting around growing large and fat.**

# WATER QUALITY

- **Good water quality means healthy happy yabbies.**
- **Good water quality means hungry fast growing yabbies. If the water quality is not good yabbies stop feeding and become susceptible to stress and pathogens.**

# FOOD

- **No food no growth.**
- **In the wild when food becomes short, the large yabbies start eating the small yabbies and the population will be reduced to match the food supply.**
- **Pond bottom needs to be covered with a layer of old hay prior to flooding with water. This hay is the main food source for yabbies.**
- **Supplemental food in the form of pellets are required to be feed 6 days per week at a rate of 0.5% biomass/day.**



# SHELTER

- **The more shelter the more yabbies you can have.**
- **Yabbies are climbing animals and will climb structures up into the water column.**
- **Caution needs to be taken with the type of shelter used**

# PREDATORS

- **Protection from predation is important. Cormorants, water rats, eels and fish are all major predators.**
- **Bird netting or protection is essential.**
- **Netting to keep the yabbies in the pond is essential.**

# POND PREPARATION

- **ADD HAY AT RATE MAX 150 BALES/Ha**
- **ADD LIME**
- **ADD WATER**

# STOCKING

- **STOCK WITH YABBIES AT 5gram MIN and 30gram MAX SIZE**
- **STOCK WITH YABBIES ALL CLOSE TO SAME SIZE**
- **STOCK AT RATE OF 2/m<sup>2</sup>**

# FEEDING

- HAY IN POND IS BASE FOOD SOURCE
- ADD EXTRA BALE EACH MONTH
- ADD PELLETS AT 0.5% (half of one %) BIOMASS/DAY - 6 DAYS/WEEK

# GROWTH

---

- **20 GRAM to 100gram in 4 – 5 MONTHS**
- **MALES GROW FASTER THAN FEMALES**
- **SINGLE SEX PONDS ARE GOOD IF POSSIBLE**

# HARVEST

- **BAITED TRAPS**
- **SPECIAL NO RING OPERA HOUSE TRAPS**
- **CATCH LARGEST FIRST**
- **BREEDING HAS OCCURRED**
- **DRAIN HARVEST**



# PRODUCTION RETURNS

- **2 CROPS/YEAR**
- **MORTALITY OF INITIAL STOCK 10 – 30%**
- **AT 30% MORTALITY & 2 CROPS = 2500Kgs/Ha/year**



# For all the information

# Buy "THE COMMERCIAL YABBY FARMER"



<b>CHAPTER 1 INTRODUCTION</b>	11
Aquaculture	12
The Yabby	15
<b>CHAPTER 2 GENERAL YABBY BIOLOGY</b>	19
Handling	19
General Identification	21
Anatomy and Sexing	22
Moulting	30
Growth	34
Breeding	36
Burrowing	42
<b>CHAPTER 3 SITE SELECTION</b>	44
Species for Aquaculture	48
<b>CHAPTER 4 YABBY HATCHERIES</b>	51
Heat	52
Air	52
Light	53
Water	53
Yabbies	54
Food and Nutrition	55
Small Tank Hatcheries	56
Still Water Hatcheries	57
Large Scale Hatcheries	60
<b>CHAPTER 5 SEMI-INTENSIVE YABBY FARMING FARM DESIGN</b>	63
Background	63
Pond Design	64
Impervious clay	65
Pond size	66
Pond depth	67
Pond bottom	67
Pond sides	67
Pond walls	68
Turbidity	69
Drainable	71
Aeration	74
Shelter	76
Overflows	80
Perimeter fence	81
Back up water supply	83
Excess Water (Effluent Pond)	84
Bird Protection	85

Pond Stocking	96
Pond Sterilization	98
<b>CHAPTER 7 FEEDING</b>	100
<b>CHAPTER 8 HARVESTING</b>	108
Draining	111
<b>CHAPTER 9 POND STOCKING/RE STOCKING</b>	113
Nursery Breeding Ponds	116
Broodstock Selection	119
Selective Breeding Programs	120
Hybrids	121
Cheran Setinas	121
Single Sex Ponds	124
<b>CHAPTER 10 PRODUCTION RETURNS</b>	126
<b>CHAPTER 11 POND MANAGEMENT</b>	128
Water Quality and Health Management	128
Temperature	129
Dissolved Oxygen	130
pH	131
Ammonia/Nitrogen	133
Salinity	134
Photosynthesis	135
Hydrogen Sulphide/Sulphur	136
<b>CHAPTER 12 PREDATORS</b>	138
Punchers - Man	138
Aquatic Predators	138
Insect Larvae	141
Eels	142
Mosquito Fish	147
Shags ( Cormorants)	148
Wading Birds	152
Kingfishers	152
Ducks and Grebes	152
Eagles and Hawks	153
Water Rats	153
Turtles	155
Others	156
<b>CHAPTER 13 DEATH AND DISEASE</b>	157
Capture/ Gill Flushing/Bacteria-	157
Disease	162
Theohamia	162
Crayfish Plague <i>Aphanomyces astaci</i>	164

Holding Cages	172
Spray Rooms	174
Holding Tanks	176
<b>CHAPTER 16 RECIRCULATION SYSTEMS</b>	180
<b>CHAPTER 17 PACKAGING</b>	194
<b>CHAPTER 18 MARKETING</b>	198
<b>CHAPTER 19 FOOD SAFETY</b>	204
<b>CHAPTER 20 COOKING</b>	209
Recipes	210
<b>CHAPTER 21 COMMERCIAL AQUA OF THE YABBY</b>	220
Design and Construction of a Commercial Farm	220
Financial Model for a Commercial Farm	222
<b>CHAPTER 22 FREQUENTLY ASKED QUESTIONS</b>	228
Yabbies	228
Fish	234
<b>CHAPTER 23 THE AUTHOR</b>	237
Conclusion	238
<b>CHAPTER 24 DEFINITIONS</b>	239
Notes	256



**Mathematics NOTES** (these may be helpful later on in the book)

1000 microns = 1 millimetre (mm)  
 1000 mm = 1 metre (mtr)  
 1 hectare = 10,000 square metres (m<sup>2</sup>)  
 1 hectare = 2.471 acres  
 1 cubic metre = 1,000 litres (ltrs)  
 1,000 litres = 1 tonne  
 1000 millilitres = 1 litre  
 1,000,000 litres = 1 megalitre (ML)  
 1,000 grams = 1 kilogram (kg)  
 1,000 kgs = 1 tonne

